

**(12) PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 199851595 B2**  
**(10) Patent No. 729454**

(54) Title  
Delivery construct for antisense nucleic acids and methods of use

(51)<sup>7</sup> International Patent Classification(s)  
C07H 021/02 C07H 021/04  
A61K 031/70 C12N 015/00

(21) Application No: 199851595

(22) Application Date: 1997.10.31

(87) WIPO No: WO98/18811

(30) Priority Data

(31) Number	(32) Date	(33) Country
08/742943	1996.10.31	US

(43) Publication Date : 1998.05.22

(43) Publication Journal Date : 1998.07.16

(44) Accepted Journal Date : 2001.02.01

(71) Applicant(s)  
The Johns Hopkins University School of Medicine

(72) Inventor(s)  
Harry C. Dietz

(74) Agent/Attorney  
F.B. RICE and CO., 139 Rathdowne Street, CARLTON VIC 3053

(56) Related Art  
PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCE USA 1996 V.93  
PAGES 7219-7224  
BIOCHEMISTRY 1994 VOL 33 PAGES 12127-12138

BEST AVAILABLE COPY

OPI DATE 22/05/98 APPLN. ID 51595/98  
AOJP. DATE 16/07/98 PCT NUMBER PCT/US97/19849



AU9851595

(PCT)

<b>(51) International Patent Classification 6 :</b> C07H 21/02, 21/04, A61K 31/70, C12N 15/00	<b>A1</b>	<b>(11) International Publication Number:</b> WO 98/18811 <b>(43) International Publication Date:</b> 7 May 1998 (07.05.98)
<b>(21) International Application Number:</b> PCT/US97/19849 <b>(22) International Filing Date:</b> 31 October 1997 (31.10.97) <b>(30) Priority Data:</b> 08/742,943 31 October 1996 (31.10.96) US <b>(71) Applicant:</b> THE JOHNS HOPKINS UNIVERSITY SCHOOL OF MEDICINE [US/US]; 720 Rutland Avenue, Baltimore, MD 21205 (US). <b>(72) Inventor:</b> DIETZ, Harry, C.; 1104 Ryegate Road, Towson, MD 21286 (US). <b>(74) Agent:</b> HAILE, Lisa, A.; Fish & Richardson P.C., Suite 1400, 4225 Executive Square, La Jolla, CA 92037 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> DELIVERY CONSTRUCT FOR ANTISENSE NUCLEIC ACIDS AND METHODS OF USE  <b>(57) Abstract</b>  A novel nucleic acid construct for delivery of antisense targeting sequences is provided. The construct includes intact stem loop structures and an antisense nucleic acid. Optionally, a ribozyme nucleic acid is included in the construct. The construct is useful for inhibition of selected genes in a cell. This allele-specific targeting is also useful in combination with replacement gene therapy.		

**DELIVERY CONSTRUCT FOR ANTISENSE NUCLEIC ACIDS**  
**AND METHODS OF USE**

5

**Field of the Invention**

This invention relates generally to the field of delivery vehicles for nucleic acid molecules and specifically to a novel construct for delivery of antisense targeting sequences to a cell.

10

**Background of the Invention**

The use of antisense oligonucleotides offers advantages over other therapeutic regimes due to their potential for target specificity. For example, conventional chemotherapy for neoplastic and virus-related diseases has the disadvantage of systemic toxicity. The therapeutic index for chemotherapeutic agents is relatively narrow, since such agents are unable to distinguish between normal and diseased cells. Antisense oligonucleotides have the potential to be many orders of magnitude more specific than traditional drugs due to their greater number of interactions with a particular target site. In theory, an oligonucleotide of more than 15-17 nucleotides in length could have the base pairing specificity to interact with only one target gene within the entire human genome. Thus, antisense oligonucleotides have the potential specificity that could serve as a powerful tool for the study of specific gene function and as therapeutic agents for disease-causing genes.

In contrast to drugs, antisense molecules are relatively simple to design. The interaction between an antisense oligonucleotide and a target mRNA is governed primarily by the sequence of the target. Oligonucleotides targeting the start codon and extending upstream or downstream have been shown to be effective. Similarly, oligonucleotides that are complementary to the splice sites have proved effective.

- 2 -

Antisense technologies for the targeted inhibition of gene expression could provide an effective strategy for the management of inherited disorders with dominant-negative or gain-of-function pathogenetic mechanisms, for the suppression of oncogenes, or for the control of a variety of infectious agents. Pathologic disorders that are currently targeted by antisense therapeutics include viral infections, inflammatory disorders, cardiovascular disease, cancers, genetic disorders and autoimmune diseases. Synthetic oligodeoxynucleotides (ODNs), especially phosphorothiates and methylphosphonates, offer the advantage of enhanced stability in biological fluids and an effectively limitless supply.

Antisense oligonucleotides are also useful for the production of transgenic animals having alterations at the germline level, such as knockout mutations, which can be used for the study of new genes or the study of the function of a known gene. Further, antisense technology combined with gene therapy is useful for example, for suppression of expression of a mutant gene product. Such gene therapy would be most advantageous in combination with a replacement regimen utilizing the "normal" gene to provide a "normal" gene product.

Unfortunately, the effective use of antisense oligonucleotides has been limited due to several problems. Disadvantages include the transient nature of ODNs, and their toxicity and propensity for producing non-sequence specified biological effects. Other disadvantages include low expression or limited stability of complementary RNAs which result in their nonspecific targeting or low efficiency of target inhibition. Antisense oligonucleotides are often poorly taken up by cells and therefore may never reach their target site. Often, antisense oligonucleotides do not reach the nucleus of a cell once administered, the site of their RNA and DNA targets. In certain applications the antisense molecules are microinjected directly into the cells. This technique works well in the laboratory, however, it cannot be applied to patients. Many of the studies with antisense show that gene expression is suppressed by 80-90% of the normal level, however, such reduction is not typically sufficient to reduce the biological effect, e.g., 10-20% expression is sufficient to maintain the biological function sought to suppress.

- 3 -

There is a need to develop a delivery system for antisense molecules that gives the antisense enhanced stability, for example by being resistant to nuclease activity or by being enriched in the nucleus, while still allowing specificity of the antisense for its target RNA or DNA. Such a system would provide effective  
5 targeting of the message with the end result being significant inhibition of expression of a particular gene.

- 4 -

### Summary of the Invention

The present invention provides a novel nucleic acid construct for delivery of antisense targeting sequences for inhibition of selected genes in a cell. The construct includes intact stem loop structures and an antisense nucleic acid. This  
5 allele-specific targeting is also useful in combination with replacement gene therapy.

In a first embodiment, the invention provides a nucleic acid construct for suppressing gene expression. The construct includes a 5' stem loop structure, an antisense nucleic acid, and a 3' stem loop structure. Such "unmodified" or "intact" stem loop structures flank the antisense nucleic acid so that the antisense  
10 oligonucleotides can readily interact with any target sequence. In a preferred embodiment, the stem loop structures are U snRNA stem loops, and most preferably, U1 snRNA stem loops. The construct provides a cloning site outside of the stem loop structures, thus rendering the stem loop structures "unmodified," into which virtually any antisense oligonucleotide could be inserted. In contrast to previous antisense  
15 delivery vehicles which have been designed to interfere with mRNA splicing, the construct of the present invention does not necessarily affect the splicing machinery so as not to disrupt normal cellular mRNA processing. In other words, while previous constructs retain elements that interact with spliceosome proteins, the construct of the invention eliminates such elements. Preferably, a U1 snRNA promoter is included in  
20 the construct. Optionally, a ribozyme nucleic acid is included in the construct.

In another embodiment, the invention provides a method for suppression of gene expression whereby a suppressive-effective amount of the nucleic acid construct of the invention is administered to a cell thereby suppressing expression of the gene. The method is particularly useful when used in combination with  
25 replacement gene therapy which provides a gene which encodes and can reconstitute the wild-type protein and which is resistant to targeting by the antisense and/or ribozyme.

The seminal discovery of the delivery vehicle of the invention now allows transgenic knockout animals to be created for the study of known or as of yet

unknown genes. For example, phenotypic consequences of loss of function due to targeted disruption of a gene(s) can be determined.

- 5 Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
22

- 6 -

**Brief Description of the Drawings**

Figure 1 shows an illustrative expression construct of the invention.

Figure 2 shows the sequence and predicted structure of chimeric transcripts derived from pU1/FIB.

5        Figure 3 shows an immunohistochemical analysis of MG63 cells that were stably transfected with an expression construct (pZeoSVLacZ, Invitrogen) lacking targeting sequence (panels A and C) or pU1/FIB (panels B and D).

10       Figure 4 shows a Northern blot analysis of 3.5 µg of poly(a) RNA extracted from confluent MG63 cells that were untransfected (lane 1), or stably transfected with either pZeoSVLacZ (Lane 2) or pU1/FIB (lane 3-5).



- 7 -

### Description of the Preferred Embodiments

The present invention provides a novel nucleic acid construct which provides a general means of delivery for antisense targeting sequences. The construct includes stabilizing structural elements, stem loop structures, which increase the effectiveness of the targeting construct. Also included is a method of suppression of gene expression utilizing the novel construct of the invention.

The nucleic acid construct of the invention which is useful for suppressing gene expression includes 5' and 3' stem loop structures and an antisense nucleic acid. While not wanting to be bound by a particular theory, it is believed that the stem loop structures provide enhanced stability of the targeting molecules by conferring resistance to exonucleases and also promote duplex formation and stability.

The "stem loop" structures refer to nucleic acid structures that have folding patterns which form hairpins and flank the antisense targeting sequence. The stem loop structures are preferably unmodified, naturally occurring structures. Alternatively, one of skill in the art would be able to synthesize such structures to "mimic" the naturally occurring structures. The preferred stem loop structures in the construct of the invention are unmodified U snRNA structures. The term "unmodified" means that the folding pattern of the stem loop structure is not compromised by alterations in the nucleic acid sequence of the naturally occurring molecule. For example, it is understood that alterations which include, but are not limited to, mutations, insertions, deletions and substitutions of one or more nucleotides can be made within the sequence of the stem loop, as long as the stabilization function and hairpin formation of the stem loop is maintained.

Preferably, the stem loop structures are derived from or synthesized according to the nucleotide sequence of the U snRNA stem loop structures. An exemplary construct utilizing U1 snRNA stem loops is shown in FIGURE 1 herein. One can produce a chimeric construct having a 5' stem loop structure from one species and a 3' structure from a second species. For example, a 5' stem loop is derived from human U snRNA and the 3' stem loop derived from a yeast U snRNA.

- 8 -

Other snRNA stem loop structures can be derived from other mammals (e.g., porcine, bovine), or yeast for example. Other chimeric constructs can be prepared using a 5' stem loop structure from one snRNA (e.g., U1) and a 3' stem loop structure from another snRNA (e.g., U3). Virtually any combination of stem loop structures is  
5 envisioned in the construct of the invention.

As used herein, the term "nucleic acid" or nucleic acid sequence" refers to a polymer of deoxyribonucleotides or ribonucleotides, in the form of a separate fragment or as a component of a larger construct. For example, nucleic acids can be assembled from cDNA fragments or from oligonucleotides to generate a synthetic  
10 gene which is capable of being expressed in a recombinant transcriptional unit. Polynucleotide or nucleic acid sequences of the invention include DNA, RNA and cDNA sequences.

Nucleic acid sequences utilized in the invention can be obtained by several methods. Sequences for specific genes or stem loop structures can be determined from  
15 published sequences and can also be found in GenBank, National Institutes of Health computer database. Nucleic acids can then be chemically synthesized by standard methods for example.

The construct of the invention includes an antisense nucleic acid flanked by the stem loop structures. The antisense nucleic acid can be directed toward any  
20 target nucleic acid, and preferably to a target message (mRNA). Any antisense which includes sequences capable of hybridizing with its complementary target can be used in the construct of the invention. For example, antisense sequences can be directed to the 5' or 3' termini of the target message, to splice junctions, or to internal sequences. One of skill in the art will readily be able to determine which sequences  
25 to use as the appropriate antisense nucleic acid construct. "Antisense nucleic acids" are DNA or RNA molecules that are complementary to at least a portion of a specific mRNA molecule (Weintraub, Scientific American, 262:40, 1990). In the cell, the antisense nucleic acids hybridize to the corresponding mRNA, forming a double-stranded molecule. The antisense nucleic acids interfere with the translation of the  
30 mRNA, since the cell will not translate a mRNA that is double-stranded. Antisense

- 9 -

oligomers of about 15 nucleotides are preferred, since they are easily synthesized and are less likely to be toxic than larger molecules when introduced into the target cell.

The use of antisense methods to inhibit the *in vitro* translation of genes is well known in the art (Marcus-Sakura, *Anal. Biochem.*, 172:289, 1988). An antisense core nucleic

5 acid may contain about 10 nucleotides complementary to the target message.

Examples of target messages include transcription regulatory factors (e.g., rent-1); viral encoded proteins (e.g., human papilloma virus E6, human immunodeficiency virus tat), (e.g., haluronic acid synthase), structural proteins (e.g., fibrillin), cytokines, oncogenes and growth factors (e.g., interleukins), etc. It should be noted that the

10 antisense nucleic acid can be useful for reducing the expression of either normal or aberrant genes.

The antisense nucleic acid can be used to block expression of a mutant protein or a dominantly active gene product, such as amyloid precursor protein that accumulates in Alzheimer's disease. Such methods are also useful for the treatment of

15 Huntington's disease, hereditary Parkinsonism, and other diseases. Antisense nucleic acids are also useful for the inhibition of expression of proteins associated with toxicity or gene products introduced into the cell, such as those introduced by an infectious agent (e.g., a virus).

It may be desirable to transfer an antisense nucleic acid encoding a

20 biological response modifier in order to reduce the expression of such biological response modifier. Included in this category are nucleic acids encoding immunopotentiating agents including a number of the cytokines classified as "interleukins". These include, for example, interleukins 1 through 12. Also included in this category, although not necessarily working according to the same mechanisms,

25 are interferons, and in particular gamma interferon ( $\gamma$ -IFN), tumor necrosis factor (TNF) and granulocyte-macrophage-colony stimulating factor (GM-CSF). It may be desirable to deliver such antisense nucleic acids to cells of the immune system to treat enzymatic related disorders or immune defects. Antisense nucleic acids can be used to reduce expression of growth factors, toxic peptides, ligands, receptors, or other

30 physiologically important proteins.

- 10 -

The construct of the invention may further include a ribozyme nucleic acid. Ribozymes are RNA molecules possessing the ability to specifically cleave other single-stranded RNA. Through the modification of nucleotide sequences which encode these RNAs, it is possible to engineer molecules that recognize specific  
5 nucleotide sequences in an RNA molecule and cleave it (Cech, *J. Amer. Med. Assn.*, 260:3030, 1988). A major advantage of this approach is that, because the ribozymes are engineered to be sequence-specific, only mRNAs with sequences complementary to the construct containing the ribozyme are inactivated.

There are two basic types of ribozymes namely, tetrahymena-type  
10 (Hasselhoff, et al., *Nature*, 334:585, 1988) and "hammerhead"-type. Tetrahymena-type ribozymes recognize sequences which are four bases in length, while "hammerhead"-type ribozymes recognize base sequences from about 3 to 18 bases in length. The longer the recognition sequence, the greater the likelihood that the sequence will occur exclusively in the target mRNA species. Consequently,  
15 hammerhead-type ribozymes are preferable to tetrahymena-type ribozymes for inactivating a specific mRNA species. The preferred ribozyme of the present invention is a hammerhead type ribozyme. The target message is required to contain a ribozyme cleavage site sequence such as 5'-GUC-3' or 5'-GUA-3' when a ribozyme is included in the construct.

20 The construct may also optionally include a triplex oligomer. Use of an oligonucleotide to stall transcription is known as the triplex strategy since the oligomer winds around double-helical DNA, forming a three-strand helix. Therefore, these triplex compounds can be designed to recognize a unique site on a chosen gene (Maher, et al., *Antisense Res. and Dev.*, 1(3):227, 1991; Helene, C., *Anticancer Drug*  
25 *Design*, 6(6):569, 1991).

The phrase "nucleic acid sequence expressing a product of interest" refers to a nucleic acid sequence which when expressed results in a product selected from a protein or antisense RNA, for example. The term "structural gene" excludes the non-coding regulatory sequence which drives transcription. The structural gene may  
30 be derived in whole or in part from any source known to the art, including a plant, a

- 11 -

fungus, an animal, a bacterial genome or episome, eukaryotic, nuclear or plasmid DNA, cDNA, viral DNA or chemically synthesized DNA. A structural gene may contain one or more modifications in either the coding or the untranslated regions which could affect the biological activity or the chemical structure of the expression  
5 product, the rate of expression or the manner of expression control. Such modifications include, but are not limited to, mutations, insertions, deletions and substitutions of one or more nucleotides. The structural gene may constitute an uninterrupted coding sequence or it may include one or more introns, bound by the appropriate splice junctions.

10           The term "operably associated" refers to functional linkage between the regulatory (e.g., promoter) sequence and the nucleic acid regulated by the regulatory sequence. The operably linked regulatory sequence controls the expression of the product expressed by the structural gene or antisense. The regulatory sequence may be homologous or heterologous to the desired gene sequence. A wide range of  
15 promoters may be utilized, including viral or mammalian promoters. Cell- or tissue-specific promoters can be utilized to target expression of gene sequences in specific cell populations. Suitable mammalian and viral promoters for the present invention are available in the art. A preferred promoter in the construct of the invention is a U snRNA promoter, and most preferably, a U1 snRNA promoter.

20           The choice of a particular heterologous promoter region as a "regulatory nucleotide sequence" of the invention, is dictated by the spatial and temporal pattern of expression that is desired for the transactivator gene and ultimately for the target transgene. Promoter regions of the invention include eukaryotically derived promoters which predominantly direct expression in, for example, the reproductive  
25 system (e.g., breast, ovary, testes); the musculoskeletal system (e.g., muscle or joint tissue); the cardiovascular system (e.g., capillaries or heart); the respiratory system (e.g., lung or nasal passages); the urological system (e.g., kidney or bladder); the gastrointestinal system (e.g., pancreas, liver, or intestines); the immune system (e.g., thymus, spleen, or circulating immunological cells); the endocrine system (e.g.,  
30 pituitary, gonads, and thyroid); the nervous system (e.g., neurons); and the

- 12 -

hematopoietic system (e.g., bone marrow and peripheral blood). Further, promoters of the invention include, but are not limited to the elastase promoter (including its enhancer; expression in pancreatic acinar cells); the alpha-A-crystallin promoter (expression in the eye lens tissue); the insulin promoter (including enhancer;  
5 expression in the pancreatic beta cells) and the albumin promoter region, including its enhancer.

Alternatively, non-eukaryotically-derived promoters such as virally-derived and prokaryotically-derived promoters are also included in the present invention. Such virally-derived promoters include, but are not limited to MMTV and  
10 MoSV LTR, SV40 early region, RSV or CMV LTR, which direct expression of viral or host genes in specific tissues and in many cell types.

Promoters useful in the invention include both constitutive and inducible natural promoters as well as engineered promoters. To be most useful, an inducible promoter should 1) provide low expression in the absence of the inducer; 2) provide  
15 high expression in the presence of the inducer; 3) use an induction scheme that does not interfere with the normal physiology of the cell; and 4) have no effect on the expression of other genes. Both constitutive and inducible promoters and enhancers will be known to those of skill in the art. The promoters used in the construct of the present invention may be modified, if desired, to affect their control characteristics.

20 The regulatory nucleotide sequence of the invention may also include "enhancer" regions. Enhancers, as used herein, refer to DNA sequences which affect transcription of a gene by RNA polymerase II, without regard to position or orientation. An enhancer region can be thousands of base pairs from a transcription unit and still affect its transcription. Example of enhancers known in the art include  
25 the immunoglobulin heavy (mu) chain or light (kappa) chain enhancers and the SV40 72-base pair repeats.

It may be desirable to deliver the construct of the invention directly to the cell without the requirement for expression, therefore, a promoter or other regulatory nucleic acid is optional. One example of a targeted delivery system for antisense  
30 polynucleotides is a colloidal dispersion system. Colloidal dispersion systems include

- 13 -

macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes or liposome formulations. The preferred colloidal system of this invention is a liposome. Liposomes are artificial membrane vesicles which are useful as delivery vehicles *in vitro* and *in vivo*. These formulations may have net cationic, anionic or neutral charge characteristics are useful characteristics with *in vitro*, *in vivo* and *ex vivo* delivery methods. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0  $\mu\text{m}$  can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley, *et al.*, *Trends Biochem. Sci.*, 6:77, 1981). In addition to mammalian cells, liposomes have been used for delivery of polynucleotides in plant, yeast and bacterial cells. In order for a liposome to be an efficient gene transfer vehicle, the following characteristics should be present: (1) encapsulation of the genes of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information (Mannino, *et al.*, *Biotechniques*, 6:682, 1988).

20           The composition of the liposome is usually a combination of p-hospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations.

25           The construct of the invention can also be delivered as a naked "gene expression vector". This means that the construct is not associated with a delivery vehicle (e.g., liposomes, colloidal particles and the like). One of the principal advantages touted for naked DNA vectors has been the lack of immune responses stimulated by the vector itself.

- 14 -

The backbone or framework of the construct preferably includes U1 snRNA nucleic acid sequences since U1 snRNA is enriched in the nucleus, is widely dispersed in the nucleoplasm, and is abundantly expressed. In the exemplary construct shown in FIGURE 1, an "antisense targeting core" was substituted for the

5 Sm protein binding site between the two naturally occurring U1 snRNA hairpins. This core contains a 30-35 basepair sequence that is directly complementary to the target message, interrupted in its center by an autocatalytic hammerhead ribozyme loop. Duplex formation aligns the ribozyme with the GUC or GUA consensus sites for ribozyme cleavage within the target message. An illustrative expression construct

10 of the invention which targets RENT1 transcripts, which are involved in the NMRD pathway, described below, is shown in Figure 1. The term *rent1* and RENT1 (regulator of nonsense transcripts) refers to the gene and protein, respectively, from either the murine or human species.

The construct described above is useful for suppressing gene expression.

15 For example, it may be desirable to modulate the expression of a gene when it is over-expressed. Where a cell proliferative disorder is associated with the expression of a gene, nucleic acid sequences that interfere with the gene's expression at the translational level can be used. This approach utilizes, for example, antisense nucleic acid, ribozymes, or triplex agents to block transcription or translation of a specific

20 mRNA, either by masking that mRNA with an antisense nucleic acid or triplex agent, or by cleaving it with a ribozyme, as described above.

The construct may also optionally include a 5' cap structure, such as a trimethylguanosine cap.

In another embodiment, the present invention also includes a method for

25 suppression of gene expression which includes administering to a cell a suppressive-effective amount of the nucleic acid construct of the invention so that the expression of the gene is suppressed. The term "suppressive-effective" amount means that amount of the construct, and thus antisense, administered is sufficient to suppress the expression of the target, *e.g.*, inhibit translation of mRNA, by at least 75% of the

30 normal expression, and preferably by at least 90%. The effectiveness of the construct



- 15 -

can be determined phenotypically or by standard Northern blot analysis or immunohistochemically, for example. Other standard nucleic acid detection techniques or alternatively immunodiagnostic techniques will be known to those of skill in the art (e.g., Western or Northwestern blot analysis).

5           The present invention also provides a method of gene therapy for the treatment of cell proliferative or immunologic disorders and diseases such as which are mediated by various proteins. The term "cell-proliferative disorder" denotes malignant as well as non-malignant cell populations which often appear to differ from the surrounding tissue both morphologically and genotypically. Such disorders may  
10 be associated, for example, with abnormal expression of a gene. "Abnormal expression" encompasses both increased or decreased levels of expression, as well as expression of a mutant form of a gene such that the normal function of the gene product is altered. Abnormal expression also includes inappropriate expression of during the cell cycle or in an incorrect cell type. The antisense polynucleotide is  
15 useful in treating malignancies of the various organ systems. Such therapy would achieve its therapeutic effect by introduction of the antisense construct into cells having the proliferative disorder. Delivery of antisense polynucleotide, can be achieved using a recombinant expression vector such as a chimeric virus or a colloidal dispersion system as described above.

20           The construct may also be useful in treating malignancies of the various organ systems, such as, for example, lung, breast, lymphoid, gastrointestinal, and genito-urinary tract as well as adenocarcinomas which include malignancies such as most colon cancers, renal-cell carcinoma, prostate cancer, leukemia, breast cancer, non-small cell carcinoma of the lung, cancer of the small intestine, and cancer of the  
25 esophagus.

          The method is also useful in treating non-malignant or immunologically-related cell-proliferative diseases such as psoriasis, pemphigus vulgaris, Bechet's syndrome, acute respiratory distress syndrome (ARDS), ischemic heart disease, post-dialysis syndrome, rheumatoid arthritis, acquired immune deficiency syndrome,  
30 vasculitis, lipid histiocytosis, septic shock and inflammation in general.

- 16 -

Various viral vectors which can be utilized for gene therapy as taught herein include adenovirus, adeno-associated virus, herpes virus, vaccinia, or, an RNA virus such as a retrovirus. Preferably, the retroviral vector is a derivative of a murine or avian retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). Preferably, when the subject is a human, a vector such as the gibbon ape leukemia virus (GaLV) is utilized. A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. By inserting a sequence of interest into the viral vector, along with another gene which encodes the ligand for a receptor on a specific target cell, for example, the vector is now target specific. Retroviral vectors can be made target specific by attaching, for example, a sugar, a glycolipid, or a protein. Preferred targeting is accomplished by using an antibody to target the retroviral vector. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome or attached to a viral envelope to allow target specific delivery of the retroviral vector containing the antisense polynucleotide.

Administration of the construct of the invention can be *in vivo*, *in vitro* or *ex vivo*.

The method of the invention also envisions gene replacement therapy to reconstitute expression of a wild-type protein in cells expressing a defective protein or, in the case of a heterozygote, a defective and a "normal" protein. Therefore, in another embodiment the invention provides a method of suppressing gene expression as above and further comprises administering a modified nucleic acid encoding a wild-type polypeptide corresponding to the gene product of the gene being suppressed, wherein the modified nucleic acid is resistant to ribozyme cleavage and/or antisense inhibition. Due to degeneracy in the genetic code, multiple nucleotide substitutions can be made across the antisense oligonucleotide target

- 17 -

region of the coding sequence within an exogenously supplied expression construct encoding the wild-type gene of interest. Such substitutions would inhibit duplex formation between the gene suppressed and the antisense nucleic acid and transcripts derived from the replacement "normal" gene, while maintaining the fidelity of the amino acid sequence. In other words, the replacement gene would provide a protein with the correct amino acid sequence utilizing codons with nucleotides that are not recognized by the antisense construct and/or ribozyme. In this method, it is therefore possible to "knockout" a defective copy of a gene/mRNA and introduce a "normal" copy resistant to the "knockout" construct. Thus, both normal and aberrant expression of a gene can be blocked by introduction of the antisense construct of the invention; to restore normal function of the gene, a copy of the wild-type gene is introduced, that, while coding for a normal protein, is resistant to duplex formation with the antisense construct and/or ribozyme cleavage.

There are a number of inherited diseases in which defective genes may be replaced including: lysosomal storage diseases such as those involving  $\beta$ -hexosaminidase or glucocerebrosidase; deficiencies in hypoxanthine phosphoribosyl transferase activity (the "Lesch-Nyhan" syndrome); amyloid polyneuropathies (-prealbumin); Duchenne's muscular dystrophy, and retinoblastoma, for example.

Pathologic disorders that can be targeted using the method of the invention include but are not limited to viral infections, inflammatory disorders, cardiovascular disease, cancers, genetic disorders and autoimmune diseases.

The method of the present invention is also useful for suppression of a dominant negative mutation. The term "dominant negative" is a term of art and refers to expression of a gene resulting in a gene product that actively interferes with the function of a "normal", endogenous protein. Thus, a mutant protein or dominantly active gene product, such as amyloid precursor protein that accumulates in Alzheimer's disease can be blocked. The dominant negative phenotype is conveyed by the expression of the mutant protein that interferes with the function of the normal protein. Such an effect is similar to the effect of dominance of one allele of a pair of alleles encoding homologous genes on a pair of homologous chromosomes such that

- 18 -

the phenotypic effect of the one allele exerts a deleterious controlling influence over the other allele.

The nucleic acid construct can be utilized in the discovery of unknown human genes, for example, with *a priori* knowledge of the gene in question. In one  
5 embodiment of this invention, cells are isolated from an individual displaying a particular phenotype which is suspected of being caused by a gene or genes containing nonsense mutations. A subset of these cells is contacted (*e.g.*, by transfecting the cells), with the construct that contains at least an antisense nucleic acid for suppression or inhibition of the expression or function of a gene (*e.g.*, Rent-1  
10 gene or other genes involved in NMRD) or its gene product in the cells. Levels of mRNA containing nonsense mutations in these cells are elevated to higher levels. Using standard techniques of differential display of mRNA, or genetic subtraction techniques commonly applied to cDNAs, the mRNA containing nonsense mutations can be enriched and then isolated by comparing the mRNA populations of the original  
15 cells with the mRNA populations in cells containing the antisense.

In a further embodiment, a transgenic animal can be developed using the novel construct and method of the invention in order to identify the impact of increased or decreased gene expression on a particular pathway or phenotype. Protocols useful in producing such transgenic animals are described below. The  
20 protocol generally follows conventional techniques for introduction of expressible transgenes into mammals. Those of ordinary skill in the art will be familiar with these applications and will be able to apply the techniques in the context of the present invention without undue experimentation.

For example, embryonic target cells at various developmental stages can  
25 be used to introduce transgenes. Different methods are used depending on the stage of development of the embryonic target cell. The zygote is the best target for microinjection. In the mouse, the male pronucleus reaches the size of approximately 20 micrometers in diameter which allows reproducible injection of 1-2pl of DNA solution. The use of zygotes as a target for gene transfer has a major advantage in  
30 that in most cases the injected DNA will be incorporated into the host gene before the

- 19 -

first cleavage (Brinster, *et al.*, *Proc. Natl. Acad. Sci. USA* 82:4438-4442, 1985). As a consequence, all cells of the transgenic non-human animal will carry the incorporated transgene. In general, this will also be reflected in the efficient transmission of the transgene to offspring of the founder since 50% of the germ cells will harbor the  
5 transgene. Microinjection of zygotes is a preferred method for incorporating transgenes in practicing the invention.

Retroviral infection can also be used to introduce transgene into a non-human animal. The developing non-human embryo can be cultured *in vitro* to the blastocyst stage. During this time, the blastomeres can be targets for retroviral  
10 infection (Jaenisch, *Proc. Natl. Acad. Sci. USA* 73:1260-1264, 1976). Efficient infection of the blastomeres is obtained by enzymatic treatment to remove the zona pellucida (Hogan, *et al.*, *Manipulating the Mouse Embryo*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). The viral vector system used to introduce the transgene is typically a replication-defective retrovirus carrying the  
15 transgene (Jahner, *et al.*, *Proc. Natl. Acad. Sci. USA*, 82:6927-6931, 1985; Van der Putten, *et al.*, *Proc. Natl. Acad. Sci. USA* 82:6148-6152). Transfection is easily and efficiently obtained by culturing the blastomeres on a monolayer of virus-producing cells (Van der Putten, *supra*; Steward, *et al.*, *EMBO J.*, 6:383-388, 1987).

Alternatively, infection can be performed at a later stage. Virus or virus-  
20 producing cells can be injected into the blastocoele (Jahner, *et al.*, *Nature*, 298:623-628, 1982). Most of the founders will be mosaic for the transgene since incorporation occurs only in a subset of the cells which formed the transgenic non-human animal. Further, the founder may contain various retroviral insertions of the transgene at different positions in the genome which generally will segregate in the offspring. In  
25 addition, it is also possible to introduce transgenes into the germ line, albeit with low efficiency, by intrauterine retroviral infection of the midgestation embryo (Jahner, *et al.*, *supra*, 1982).

A third type of target cell for introduction of heterologous nucleic acid sequences is the embryonal stem cell (ES). ES cells are obtained from pre-  
30 implantation embryos cultured *in vitro* and fused with embryos (Evans, *et al.*, *Nature*,

- 20 -

292:154-156, 1981; Bradley, *et al.*, *Nature*, 309:255-258, 1984; Gossler, *et al.*, *Proc. Natl. Acad. Sci USA*, 83:9065-9069, 1986; and Robertson, *et al.*, *Nature*, 322:445-448, 1986). Transgenes can be efficiently introduced into the ES cells by DNA transfection or by retrovirus-mediated transduction. These transformed ES cells can thereafter be combined with blastocysts from a non-human animal. The ES cells will thereafter colonize the embryo and contribute to the germ line of the resulting chimeric animal (see for review, Jaenisch, *Science*, 240:1468-1474, 1988). Any ES cell may be used in accordance with the present invention. It is, however, preferred to use primary isolates of ES cells. Such isolates may be obtained directly from embryos such as with the CCE cell line disclosed by Robertson, E.J., in *Current Communications in Molecular biology*, Capecchi, M.R. (Ed.) Cold Springs Harbor Press, Cold Springs Harbor, NY (1989), pp.39-44), or from the clonal isolation of ES cells from the CCE cell line (Schwartzberg, P.A. *et al.*, *Science* 246:799 (1989). ES cells may be derived or isolated from any species, although cells derived or isolated from mammals such as rodents, rabbits, and non-human primates are preferred.

Construction of transgenes can be performed by those of ordinary skill in the art using the teachings herein. One of ordinary skill in the art can "knock out" a gene in mice by targeted disruption of the gene. This can be accomplished by homologous recombination in murine embryonic stem (ES) cells using standard techniques. The clinical and cellular consequences of targeted disruption can be investigated in multiple genetic backgrounds including inbred strains, strains with many undefined nonsense alleles, and strains of known mutant genotype to determine if: (a) targeted disruption of the gene can effect a loss of function; (b) loss of function can have an phenotype consequences (i.e., the creation of knockout phenotypes), (c) loss of the gene product unmasks the effects of protein that can be expressed from "physiologic" somatically acquired or inherited nonsense alleles upon transcript stabilization.

The cre/lox system as described in US Patent Serial No. 4,959,317, incorporated herein by reference, can be utilized in the production of transgenic animals. A first and second loxP DNA sequence is introduced into cells connected by

- 21 -

- a pre-selected antisense or replacement gene, herein referred to as a "target transgene". The "target transgene" of interest can be a complete gene or any other sequence of nucleotides including those of homologous, heterologous, or synthetic origin. The target transgene sequence can be for example, an antisense or
- 5 replacement gene for a structural protein, an enzyme, or a regulatory molecule. The target transgene may also be a gene of undetermined function. Using tissue-specific or developmentally-specific regulatory sequences (as described above) to direct expression of the target transgene, a function could be identified. If the first and second lox sites have the same orientation (direct repeats), activation of the regulatory
- 10 nucleotide sequence of the transactivator transgene results in a deletion of the target transgene DNA, such that ablation or modification of activity results. If the first and second lox sites have opposite orientation (inverted repeats), activation of the regulatory nucleotide produces an inversion of the nucleotides sequence of the target transgene.
- 15 The construct of the invention may be used to introduce DNA sequences into the germ line cells of "non-humans" to create transgenic animals. The preferred animal of the invention is a mouse. However, other non-humans of the invention include but are not limited to other rodents (e.g. rat, hamster), rabbits, chickens, sheep, goats, fish, pigs, cattle, and non-human primates.
- 20 In yet another embodiment, an antisense library can be constructed using the nucleic acid construct of the invention in order to identify novel genes. For example, a UI construct library containing antisense core sequences (e.g., random sequences or sequences derived from a cDNA library) is transfected into host cells and clones exhibiting a desired phenotype are identified. Such clones are then
- 25 examined to determine the nature of the regulatory sequence in the construct. Therefore, novel genes having specific functions can be identified. For example, cells that acquire a transformed phenotype may contain an antisense molecule for a novel tumor suppressor gene. A match between a given suppressive nucleic acid and a previously cloned gene can be made using standard techniques (e.g., BLAST search
- 30 algorithm).

- 22 -

The following Examples are intended to illustrate, but not to limit the invention. While such Examples are typical of those that might be used, other procedures known to those skilled in the art may alternatively be utilized.



- 23 -

### EXAMPLES

The following examples provide a description of construction of an exemplary nucleic acid construct of the invention, utilizing 5' and 3' unmodified stem loop structures from U1 snRNA which flank an antisense molecule directed toward the fibrillin-1 gene (mRNA).

#### EXAMPLE 1

An antisense expression construct was constructed that incorporates several potentially enhancing features. The pU1/FIB vector was constructed on the backbone of the pZeoSV (Invitrogen) prokaryotic/eukaryotic expression vector. The SV40 promoter, polyadenylation site and polylinker were excised from pZeoSV at the BamHI sites. A U1 snRNA expression cassette cloned into pUC13 was excised with BamHI digestion and ligated into the BamHI sites of the modified pZeoSV. Two rounds of site-directed mutagenesis (Deng, et al., *Anal. Biochem.*, 200, 81; 1992) were then performed to change four nucleotides flanking the Sm protein restriction sites (pZeoU1EcoSpe). Complementary oligonucleotides that encode the antisense 'core' sequence, shown in Figure 1, including the 24 highly conserved nucleotides of hammerhead ribozymes (Cech, et al., *Ann. Rev. Biochem.*, 55:599, were synthesized and annealed at 40° C such that the remaining 5' and 3' overhangs were exactly complementary to the overhangs left by EcoRI and SpeI digestion. The sequences of the oligonucleotides were as follows:

5'-AATTGGCGATCTCCAGCACTGATGAGTCCGTGAGG  
ACGAAACGCCCTCGACGCAT-3' (SEQ ID NO:1),  
5'-CTAGATGCGTCGAGGGCGTTTCGTCCTCACGGACTC  
ATCAGTGCTGGAGATCGCC-3' (SEQ ID NO:2) (sense and antisense,  
respectively). The resulting duplex was ligated into the EcoRI and SpeI sites of pZeoU1EcoSpe to create pU1FIB. All ligation junctions were sequenced to verify the identity and orientation of the insert.

- 24 -

In that the structure described for naturally occurring antisense RNAs is highly similar to that for small nuclear RNAs (snRNAs), essential components of the spliceosome complex that are abundant and stable in the nucleus of mammalian cells, the U1 snRNA gene was selected as the framework for vector construction (Guthrie, et al., *Annu. Rev. Genet.*, 22; 387, 1988). Other attributes include the potent and constitutively active nature of the U1 snRNA promoter, the ability of the unusual trimethylguanosine 5' cap and Sm protein interactions to signal transport of U1 snRNA back into the nucleus (Hamm, et al., *Cell*, 62; 569, 1990; Fischer, et al., *Science* 249; 786, 1990; Fischer, et al., *J. Cell Biol.* 113 Mo; 705, 1991; Plessel, et al., *Mol. Cell. Biol.* 14, 4160, 1994), and the lack of polyadenylation of mature snRNAs, a factor which may favorably influence transcript trafficking and localization (Zhong, et al., *Proc. Natl. Acad. Sci. USA*, 91; 4258, 1994). Moreover, unlike other spliceosome components, U1 snRNA is widely dispersed in the nucleoplasm (Carmo-Fonseca, et al., *EMBO J.*, 10; 195, 1991; Matera, et al., *J. Cell. Biol.*, 121; 715, 1993).

As an illustrative example, the antisense targeting 'core' was designed to contain sequences exactly complementary to coding nucleotides 1-15 and 17-30 of fibrillin mRNA (Corson, et al., *Genomics*, 17; 476, 1993; Pereira, et al., *Hum. Mol. Genet.*, 2; 961, 1993) separated by the 22 bp hammerhead ribozyme loop. Dominant-negative forms of fibrillin-1 cause Marfan syndrome, an autosomal dominant systemic disorder of connective tissue (Dietz, et al., *Nature*, 352; 337, 1991). The regions of complementarity were predicted to align the autocatalytic structure with the consensus sequence for ribozyme cleavage (5'-GUC-3') within the target message (Cech, et al., *Annu. Rev. Biochem.*, 55; 599, 1986; T.R. Cech, *Science*, 236; 1532, 1987; T.R. Cech, *Ann. Rev. Biochem.*, 59, 543, 1990; Haseloff, et al., *Nature*, 334; 585, 1988). This antisense targeting sequence was substituted for the short Sm protein binding sequence between the two hairpin loops of U1 snRNA (Guthrie, *Annu. Rev. Genet.*, 22; 387, 1988.) The sequence of the resulting chimeric RNA (Figure 1) was analyzed using a program that predicts RNA structure (Mike Zuker's RNA page: <http://ibc.wustl.edu/~zucker/rna/>). The boundaries of the targeting sequence utilized in the construct were selected to maximize preservation of the U1

- 25 -

snRNA stem-loops, the ribozyme secondary structure and the accessibility of the sequence complementary to the target message. Expression of the chimeric RNA was under the transcriptional control of a region of U1 snRNA 5'-flanking sequence that has been shown to be potent and constitutively active (Zhuang, et al., *Cell*, 46; 827, 1986; Maugin, et al., *EMBO J.*, 5; 987, 1986); and Asselbeyers, et al., *Mol. Biol. Rep.* 17; 101, 1993).

Figure 2 shows the sequence and predicted structure of chimeric transcripts derived from pU1/FIB. The Sm protein binding site of U1 snRNA (boxed) has been substituted by an antisense targeting core (flanked by dashed lines) that is complementary to the first 30 coding nucleotides of FBN1 and contains a hammerhead ribozyme sequence in its center. The two stem-loop structures of U1 snRNA are maintained. Cleavage of the target message (unbolded characters) is predicted to occur immediately following the consensus sequence 5'-GUC-3' (arrow). The 5' cap structure of the chimeric targeting molecule may contain 1 or 3 methyl (m) groups. A sequence in the chimeric molecule (5'-AAUUGG-3', underlined) remains highly similar to the consensus for Sm protein binding, PuA(U)nGPu (Pu = A or G).

A human osteosarcoma (MG63) cell line was stably transfected with either a reporter gene expression vector (pZeoSVLacZ, Invitrogen) or the chimeric construct (pU1/FIB) by long-term selection for zeomycin resistance. Human osteosarcoma (MG63) cells (Sakai, et al., *J. Cell Biol.*, 103; 2499, 1986) were grown to 60% confluency and transfected with either linearized pU1/FIB or a lac Z reporter gene construct (pZeoSVLacZ, Invitrogen), used as a control. The transfections were performed using a DOTAP liposome formulation (Boehringer Mannheim, 1 mg/ml) according to manufacturer's instructions. Cells were grown in MEM media (Cellgro) with 10% FCS and 250 µg/ml of zeocin (Invitrogen), and were maintained in selection for 14 days prior to the isolation of clonal colonies. Mono- and polyclonal colonies were established and grown to confluency.

Figure 3 shows an immunohistochemical analysis of MG63 cells that were stably transfected with an expression construct (pZeoSVLacZ, Invitrogen) lacking targeting sequence (panels A and C) or pU1/FIB (panels B and D). The antibodies

- 26 -

either recognized epitopes in fibrillin-1 (A and B) or fibronectin (C and D). Analysis was performed in duplicate for each of three independent clonal colonies for each construct, with identical results. Representative fields are shown at 63X magnification. Untransfected MG63 cells showed a pattern of protein deposition that  
5 was indistinguishable from that for cells transfected with pZeoSVLacZ.

Fibrillin-1 was strikingly absent upon immunohistochemical analysis (Eldadah et al., *J. Clin. Invest.*, 95; 874, 1995) using either anti-fibrillin-1 mAb 69 (a gift from L. Y. Sakai) or an anti-fibronectin mAb (Sigma) of cells harboring pU1/FIB, while cells transfected with pZeoSVLacZ showed a pattern of protein  
10 deposition that was indistinguishable from untransfected controls (Figure 3). The lines harboring pU1/FIB could not be distinguished from mock-transfected or untransfected cells upon immunohistochemical analysis with a monoclonal antibody to fibronectin, suggesting specificity for the targeting process.

Antisense RNAs expressed by pU1/FIB might inhibit fibrillin-1 expression  
15 by multiple mechanisms. Ribozyme cleavage would remove the 5' cap from targeted transcripts, an event predicted to effect their rapid degradation. Alternatively, in the absence of ribozyme cleavage, involvement of the initiating AUG codon in duplex formation would impair translation.

The magnetic porous glass direct mRNA purification technique was used  
20 to isolate poly(A) RNA (according to the manufacturer's instructions, CPG, Inc.) and Northern blot analyses were performed. Electrophoresis of 3.5 µg of mRNA was performed under denaturing conditions, as previously described (Lehrach, et al., *Biochemistry*, 16; 4743, 1977; D.A. Goldberg, *Proc. Natl. Acad. Sci. USA*, 77; 5794, 1980). The gel was exposed to 60 mJoules UV light to facilitate transfer of large  
25 mRNA species. RNA was transferred to nylon membrane using the turboblotting system according to manufacturer's instructions (Schleicher & Schuell). The membrane was washed briefly in 2X SSC, crosslinked with 125 mJoules of UV light and prehybridized in Expresshyb (Clontech) for 1/2 hour at 68° C. Human cDNA probes encoding fibrillin-1 (nt 370-1183), β-actin (Clontech), and G3PDH (Clontech)  
30 were labeled by random priming (Feinberg, et al., *Anal. Biochem.*, 132; 6, 1983). The

- 27 -

membrane was hybridized for 1 hour at 68° C, washed first in 2X SSC with 0.05% SDS at room temperature and then in 0.1X SSC with 0.1% SDS at 50° C.

Figure 4 shows a Northern blot analysis of 3.5 µg of poly(a) RNA extracted from confluent MG63 cells that were untransfected (lane 1), or stably  
5 transfected with either pZeoSVLacZ (Lane 2) or pU1/FIB (lane 3-5). Cells for lanes 1 and 3 were polyclonal, while cells for lanes 2, 4 and 5 were derived from monoclonal colonies. Hybridization was performed with radiolabeled cDNA probes complementary to fibrillin-1, β-actin, and G3PDH transcripts.

Northern blot analysis of multiple clonal colonies harboring pU1/FIB  
10 revealed undetectable levels of FBN1 message (Figure 4). FBN1 transcripts were easily detected upon Northern analysis of RNA extracted from untransfected and mock-transfected cells. All cell types showed comparable amounts of β-actin and G3PDH RNA, confirming the specificity of targeting.

A similar U1 construct was created containing U1 snRNA stemloop  
15 structures and an antisense core aimed at inhibiting rent-1 expression. Functional analysis of transfected cells showed a significant up-regulation of the steady-state abundance of transcripts derived from endogenous nonsense alleles.

### SUMMARY

These data suggest that most, if not all of the inhibitory effect was  
20 achieved at the level of target message abundance, presumably by reducing mRNA stability. Although an influence upon transcriptional efficiency cannot be excluded, it is not intuitive given the site and nature of the predicted targeting sequence-target interaction. Based upon previous studies, the antisense-induced mRNA degradation most likely takes place within the nucleus (Murray, et al., in *Modern Cell Biology*,  
25 J.A.H. Murray, Ed. (Wiley/Liss, New York, 1992), pp 1-49 M.Comelissen, *Nucleic Acids Res.*, 17; 7203, 1989.). Three elements of nuclear pre-U1 RNA, the 5' m7G cap, the 3' terminal stem loop, and sequences in the 5'-terminal 124 nucleotides, contribute to the stability of pre-U1 and snRNA and to efficient nuclear export (Neuman de Vegvar, et al., *Mol. Cell. Biol.*, 10; 3365, 1990; Tems, et al., *Genes Dev.*,

- 28 -

- 7; 1898, 1993; Yuo, et al., *ibid.*, 3; 697, 1989; Izaurralde, et al., *Nature*, 376; 709, 1995). Nuclear targeting of cytoplasmic U1 snRNA is influenced by hypermethylation of the 5' cap structure and binding of at least one common U snRNP protein, both dependent upon the integrity of the Sm protein binding site which has
- 5 been altered in pU1/FIB (Guthrie, et al., *Annu. Rev. Genet.*, 22; 387, 1988). A sequence at the 5' end of the insert (AAUUGG) is highly similar to the consensus site [PuA(U)nGPu, Pu = A or G] for Sm protein binding (Jones, et al., *EMBO J.*, 9; 2555, 1990). It has also been shown that Sm binding sites are highly tolerant of mutations including internal nucleotide substitutions and deletions (Jones, et al., *supra*).
- 10 Stable transfection of cultured cells with an expression vector containing the isolated FBN1/ribozyme core sequence of pU1/FIB but no U1 snRNA secondary structure, under the transcriptional control of the SV40 promoter, resulted in no discernible inhibitory effect on the expression of fibrillin-1 transcript or protein. The oligonucleotides described above containing the 24 highly conserved nucleotides of
- 15 hammerhead ribozymes flanked by complementary sequence for 10 codons in the translation start site region of the FBN1 mRNA, were cloned into the SpeI and EcoRI restriction sites for the polylinker of pZeoSV (Invitrogen). Fetal fibroblasts were stably transfected with this vector. Immunohistochemical analysis of these clones was performed by methods identical to those described herein). These results suggest
- 20 that selected properties of U1 snRNA and/or its promoter are contributing to the extreme efficiency of inhibition seen with the use of the chimeric cRNA.

Although the invention has been described with reference to the presently preferred embodiment, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention

25 is limited only by the following claims.

- 29 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: The Johns Hopkins University School of Medicine
- (ii) TITLE OF INVENTION: Delivery Construct for Antisense Nucleic Acids and Methods of Use
- (iii) NUMBER OF SEQUENCES: 2
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Fish & Richardson P.C.
  - (B) STREET: 4225 Executive Square, Suite 1400
  - (C) CITY: La Jolla
  - (D) STATE: CA
  - (E) COUNTRY: USA
  - (F) ZIP: 92037
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: 08/742,943
  - (B) FILING DATE: October 31, 1996
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Haile, Lisa A.
  - (B) REGISTRATION NUMBER: 38,347
  - (C) REFERENCE/DOCKET NUMBER: 07265/098WO1
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: 619/678-5070
  - (B) TELEFAX: 619/678-5099

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 55 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

- 30 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AATTGGCGAT CTCCAGCACT GATGAGTCCG TGAGGACGAA ACGCCCTCGA CGCAT 55

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS: --

- (A) LENGTH: 55 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CTAGATGCGT CGAGGGCGTT TCGTCCTCAC GGACTCATCA GTGCTGGAGA TCGCC 55



## THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A nucleic acid construct for suppressing expression of a target gene, comprising:
  - 5 an antisense nucleic acid sequence directed to a target gene of interest;
  - 5' of said antisense nucleic acid region, an unmodified, natural occurring 5' U snRNA stem loop structure;
  - a pol II promoter region 5' of said antisense nucleic acid sequence;
  - 10 and
  - 3' of said antisense region, an unmodified, naturally occurring 3' U snRNA stem loop structure,
  - wherein said expression of said target gene is suppressed by at least 75% of the normal level of expression.
- 15 2. The nucleic acid construct of claim 1, wherein the U snRNA is U1.
3. The nucleic acid construct of claim 1 or claim 2 wherein the promoter is a U1 snRNA promoter.
- 20 4. The nucleic acid construct of claim 1 or claim 2 wherein the promoter is a constitutive promoter.
5. The nucleic acid construct of claim 1 or claim 2 wherein the promoter is an inducible promoter.
- 25 6. The nucleic acid construct according to any one of claims 1 to 5, further comprising a ribozyme nucleic acid.
- 30 7. The nucleic acid construct of claim 6, wherein the ribozyme nucleic acid is located between the 5' and 3' stem loop structures.
8. The nucleic acid construct of claim 6 or claim 7 wherein the ribozyme nucleic acid is a hammerhead-type ribozyme.



9. The nucleic acid construct of claim 6 or claim 7 wherein a consensus sequence for ribozyme cleavage in a target nucleic acid is 5'-GUC-3' or 5'-GUA-3'.

5

10. The nucleic acid construct according to any one of claims 1 to 9, wherein the antisense nucleic acid is targeted to a region of a gene is selected from the group consisting of rent-1, HPV E6, HIV, hyaluronic acid synthase, and fibrillin.

10

11. A method for suppression of gene expression comprising administering to a cell a suppressive-effective amount of the nucleic acid construct according to any one of claims 1 to 10.

15

12. The method of claim 11, wherein the administering is *in vitro*.

13. The method of claim 11, wherein the administering is *in vivo*.

14. The method of claim 11, wherein the administering is *ex vivo*.

20

15. The method according to any one of claims 11 to 14, further comprising:  
administering a second nucleic acid encoding a wild-type polypeptide corresponding to the gene product of the gene being suppressed, wherein the second nucleic acid is resistant to ribozyme cleavage and/or antisense inhibition.

25

16. A non-human transgenic animal comprising the nucleic acid construct according to any one of claims 1 to 10.

30

17. A non-human knock-out animal produced utilizing the nucleic acid construct according to any one of claims 1 to 10.

18. The nucleic acid construct according to any one of claims 1 to 10, further comprising a 5' trimethylguanosine cap.



Dated this fifteenth day of November 2000

The Johns Hopkins University School  
of Medicine

Patent Attorneys for the Applicant:


F B RICE & CO

850 551 10





FIG. 2



Plasmid name: pU1/RIBOZYME


Plasmid size: 3610 bp

Constructed by: H. C. Dietz & R. A. Montgomery

Construction date: 11/20/95


Comments/References: none

FIG. 3



BEST AVAILABLE COPY

FIG. 4



BEST AVAILABLE COPY

BEST AVAILABLE COPY